

AD-A052 410

CALIFORNIA UNIV SANTA BARBARA QUANTUM INST
DEVELOPMENTAL STUDIES OF POTENTIAL CHEMICALLY PUMPED ELECTRONIC--ETC(U)
MAR 78 D J BENARD, W D SLAFER, P H LEE

F/G 20/5

AFOSR-76-2959

AFOSR-TR-78-0693

NL

UNCLASSIFIED

| OF |
AD
A052410



END
DATE
FILMED
5-78

DDC



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR TR- 78 - 0693	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DEVELOPMENTAL STUDIES OF POTENTIAL CHEMICALLY PUMPED ELECTRONIC TRANSITION LASERS: COMBUSTION OF ALKALINE EARTHS IN NITROUS OXIDE-CARBON MONOXIDE AND REACTIONS OF METASTABLE ALKALINE EARTH ATOMS.		5. TYPE OF REPORT & PERIOD COVERED Final Scientific Report 10/15/75 - 10/14/77
6. PERFORMING ORG. REPORT NUMBER		7. CONTRACT OR GRANT NUMBER(s) AFOSR-76-2959
8. AUTHOR(s) David J. Benard, Warren D. Slafer Paul H. Lee		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2303/B1
9. PERFORMING ORGANIZATION NAME AND ADDRESS Quantum Institute University of California Santa Barbara, CA 93106		10. REPORT DATE March 10, 1978 10 Mar 78
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (AFSC) Bolling Air Force Base, D.C. 20332		12. NUMBER OF PAGES 15 15 p.
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Final scientific rept. 15 Oct 75 - 14 Oct 77		14. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES In addition to 16 copies of the Final Report, enclosed is: 1) one set of 30 figures used in presentations, reports, and publications throughout the grant period; 2) one set of all reprints & pre-prints submitted under grant.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) chemical laser, metastable alkaline earth atoms, KMg, metal oxidation flames, gain measurement, metal hydride, energy transfer		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Alkaline earth catalyzed N ₂ O-CO flames have been studied as potential electronic transition chemical lasers. The addition of CO to the M+N ₂ O flames, where M=Mg, Ca, and Sr, has been shown to efficiently produce the respective metastable alkaline earth (P) atoms. These flames have been further shown to exhibit enhanced MO(A-X) photon yields as well as new MO band systems in the chemiluminescence spectra. The mechanism of these reactions has been elucidated and information has been gained regarding -continued over-		

407 624

alt

DDC FILE COPY 9 AD A 052410

DDC
APR 10 1978
F

Block 20. - continued

the identity of the reservoir state. Results of spectroscopic assignments and gain measurements on the alkaline earth oxide bands are reported as are results of transfer and excimer formation studies.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
D. C.	SPECIAL
A	

Unclassified

AFOSR-TR- 78 - 0693

FINAL SCIENTIFIC REPORT

DEVELOPMENTAL STUDIES OF POTENTIAL CHEMICALLY PUMPED
ELECTRONIC TRANSITION LASERS:
COMBUSTION OF ALKALINE EARTHS IN N_2O+CO
AND REACTIONS OF METASTABLE ALKALINE EARTH ATOMS

David J. Benard
Warren D. Slafer
Paul H. Lee

Quantum Institute
University of California
Santa Barbara, California 93106

Approved for public release;
distribution unlimited.

Grant: AFOSR-76-2959
Period: October 15, 1975 - October 14, 1977
Award: \$102,812
Date: March 10, 1978

prepared for

Air Force Office of Scientific Research (AFSC)
Directorate of Chemical Sciences
Bolling Air Force Base, D.C. 20332

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSO)

NOTICE OF TRANSMITTAL TO DDC

This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.

A. D. BLOSE

Technical Information Officer

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	1
II. FINDINGS	1
1. M-N ₂ O-CO Flames	1
2. Mechanism of Reaction	2
3. The Reservoir State	3
4. Reactions of M(³ P)	4
5. Spectroscopic Assignments	4
6. Gain Measurements	5
7. The MO Ground State	5
8. Energy Transfer Studies	6
9. Metal Hydrides	7
10. Bimetallic Molecules	7
11. Bound-Free Transitions	7
III. RECOMMENDATIONS	8
IV. REFERENCES	11
APPENDIX	12

I. INTRODUCTION

This report deals with a two year study of alkaline earth catalyzed N_2O -CO flames as potential electronic transition chemical lasers. The details of the experiments and the analysis of the results have all been published in the open literature, therefore, we summarize herein only the significant findings and make recommendations as to future efforts.

II. FINDINGS

Metal oxidation flames such as $Ba+N_2O$ have been intensively studied as laser candidates. Central to these studies lies the concept of a long lived or metastable reservoir state that is the adiabatic reactions product.⁽¹⁾ Many of the results that we have obtained deal with this state, which was previously sensed by us in CO_2 energy transfer experiments.⁽²⁾ These early experiments provided evidence for efficient energy storage in chemically generated MgO ($Mg+N_2O$ reaction) and for chemical reaction of the MgO reservoir state with CO. These findings set the background for our initial investigations of $M-N_2O$ -CO flames.


1. $M-N_2O$ -CO Flames

Adding CO to a $M+N_2O$ flame, where $M=Mg$, Ca and Sr, had several interesting effects, including (1) efficient production of $M(^3P)$ metastable species, (2) enhanced $MO(A\rightarrow X)$ photon yields

and (3) the appearance of new MO band systems in the chemiluminescence spectra. The production of $M(^3P)$ species appeared to be essentially 100% efficient; however, the metastables once generated were, of course, subject to quenching and further reaction. The MO photon yields were increased by as much as a factor of 40 (MgO, B→X system) upon addition of CO and in this way an absolute SrO, A→X photon yield of 22% was obtained.

2. Mechanism of Reaction

The gases N_2O and CO do not react rapidly at low pressure and temperature unless catalyzed.⁽³⁾ Fennimore⁽⁴⁾ found that catalysis by Na was rapid and resulted in efficient chemi-excitation of the Na atoms to the first excited p-state. We found similar results for catalysis by Mg, Ca and Sr but not for Ba. The mechanism appears to be



where MO^+ is the metal oxide reservoir state, since the second step is an endothermic reaction for ground state MO to produce M^* species. Coproduction of CO_2 with M^* was also demonstrated using CO_2 laser induced fluorescence techniques.

3. The Reservoir State

The identity of the MO reservoir state was of much interest. Our results show that this state is much more stable in the lighter MO species. This result can be understood simply in terms of the energy overlap of the MO emitting states with the heat of reaction; whereas the $\text{Mg-N}_2\text{O}$ reaction exoergicity fails to reach the $\text{MgO}(\text{B}^1\Sigma)$ state, the $\text{Ba+N}_2\text{O}$ heat release is in good resonance with the $\text{BaO } a^3\Pi, ^1\Pi$ states and the emitting $\text{A}^1\Sigma$ state. Thus BaO chemiluminescence is very bright in comparison to MgO, but the MgO retains its chemical excitation over a much longer time scale. The efficiency of the M^* generation suggests that the $\text{M+N}_2\text{O}$ reaction produces the MO^+ reservoir state with an efficiency that is essentially independent of molecular weight. This means that laser schemes can profitably be based on Mg flames as well as Ba systems, thus allowing one to take advantage of the superior volatilization properties of the lighter metal.

The identity of the reservoir state was sought after and initial quenching experiments tended to indicate a vibrationally excited species. We found that the continuum emitted by $\text{M+N}_2\text{O}$ flames was not the chemiluminescence of the MO^+ reservoir state, even though it was quenched upon addition of CO, but rather that the continuum derived its energy from the MO^+ state. The carrier of this feature remains unknown but may be polyatomic,

such as $(\text{MO})_2$, which would explain the lack of spectroscopic structure and the fact that energy pooling must be involved, since the continuum extends to photon energies well beyond the $\text{M}+\text{N}_2\text{O}$ exoergicity.

4. Reactions of $\text{M}(^3\text{P})$

The primary end of the chemically generated $\text{M}(^3\text{P})$ in our flames was to react again with N_2O and produce highly excited MO species. In the case of Mg flames, the $\text{Mg}^*+\text{N}_2\text{O}$ reaction was shown to lead by correlation rules to the $\text{MgO}(^3\Delta)$ state, which decays to the $\text{a}^3\Pi$ state in a narrow band system centered at 372 nm. The $\text{Mg}^*+\text{N}_2\text{O}$ reaction was also found to populate the B state as well, which is not expected if correlation arguments hold. This may, however, be the result of a non-adiabatic process, since the $\text{Mg}^*+\text{N}_2\text{O}$ reaction may also go adiabatically to $\text{MgO}(\text{a}^3\Pi)$ with sufficient vibrational excitation to transfer into the B state upon subsequent collisions.

5. Spectroscopic Assignments

The results with Ca and Sr were analogous to those with Mg. The bands of CaO and SrO, known as "arc bands", were excited by reactions of Ca^* or Sr^* with N_2O in flames containing CO. The assignment of the "arc bands" has been largely confused with "flame bands" due to MOH.⁽⁵⁾ We established that the systems present in the $\text{M}+\text{N}_2\text{O}+\text{CO}$ flames are not MOH

nor are they $(\text{MO})_2$, as though by Gaydon.⁽⁶⁾ Eckstrom⁽⁷⁾ and Field⁽⁸⁾ have also studied the 550 nm band system we assigned to CaO ($^3\Delta \rightarrow a^3\Pi$) and concur in our findings. The CaO and SrO bands are analogous to the known ultraviolet band systems of MgO .⁽⁹⁾

6. Gain Measurements

Several diagnostic measurements were employed to assay for gain on any of the chemiluminescent transitions inherent in the $\text{M-N}_2\text{O-CO}$ flames. We found that the $\text{M}(^3\text{P}_1 \rightarrow ^1\text{S}_0)$ and $\text{MO}(^3\Delta, ^1\Delta + ^1\Sigma^- \rightarrow a^3\Pi, ^1\Pi)$ transitions were not inverted. However, as a result of these efforts we did find very large populations in the CaO and SrO $a^3\Pi, ^1\Pi$ states and we further showed, based on the way in which these concentrations varied with CO , that the $a^3\Pi, ^1\Pi$ states were not the MO^+ reservoir or energy pooling states.

7. The MO Ground State

Harris, Revelli and Wicke⁽¹⁰⁾ have reported anomalously low $\text{BaO}(X^1\Sigma)$ concentrations in $\text{Ba+N}_2\text{O}$ flames. Using an absorption probe we obtained similar results for $\text{Ca+N}_2\text{O}$ and $\text{Sr+N}_2\text{O}$ and confirmed the results for the $\text{Ba+N}_2\text{O}$ flames. The absorptions on the $\text{MO}(A \rightarrow X)$ transitions were two orders of magnitude lower than those recorded for the $\text{MO}(^3\Delta, ^1\Delta + ^1\Sigma^- \rightarrow a^3\Pi, ^1\Pi)$ transitions in identical flames. These results imply a rapid removal mechanism for the $\text{MO}(X^1\Sigma)$ ground state and a probable

inversion of the $\text{MO}(a^3\Pi, ^1\Pi \rightarrow X^1\Sigma)$ band systems. These forbidden transitions are, however, too weak to produce a laser based on a molecular species. Adding CO to the $\text{M}+\text{N}_2\text{O}$ flames decreased still further the $\text{MO}(X^1\Sigma)$ ground state while initially enhancing (at least for CaO and SrO) the concentration of the A states. Had this trend continued, which was not the case, inversion of the strong chemiluminescent $A \rightarrow X$ transitions could have been obtained.

As a result of these studies, we concluded that there are no useful inversions inherent in the $\text{M}+\text{N}_2\text{O}+\text{CO}$ flames and that laser potential, if any, in such flame systems lies with their use as $\text{M}(^3\text{P})$ generators.

8. Energy Transfer Studies


We attempted briefly and without success to observe energy transfer from chemi-excited MgO to iodine atoms. Much greater success was achieved by introducing Ca into a flowing after-glow of $\text{Mg}(^3\text{P})$ atoms. The Mg^* was rapidly quenched and the Ca atoms were excited, despite the lack of resonance between atomic energy levels. The Mg-Ca transfer system was further studied by electrical discharge in a mixed metal heat pipe. Evidence was obtained for the pumping of Ca by Mg^* , but deleterious energy pooling reactions (Mg^*+Ca^*) were responsible for preventing inversion.

9. Metal Hydrides

We also observed very intense MgH and CaH emission in RF discharged metal vapor heat pipes where trace impurities of water vapor had contaminated the sample. Electrical-to-optical conversion efficiencies as high as 5% were measured. The probable mechanism was excitation of the metals to the 3P state by the discharge, which is a very efficient process, followed by resonant energy transfer to the metal hydrides which then decay by a set of highly allowed transitions to the ground state. The potential curves of MgH or CaH are, however, such as to discourage any possibility of inversion.

10. Bimetallic Molecules

We also attempted to observe energy transfer from Mg(3P) to K atoms in the afterglow. The nearly resonant K(5s) state was not excited although the K(4p) energy level was. The molecule KMg was, however, observed to form rapidly in the excited state with efficiencies of up to 20%. Subsequent investigations showed that the pumping reaction was



11. Bound-Free Transitions

It was speculated that KMg might be an excimer and that the KMg chemiluminescence might consist of bound-free transitions.

A laser induced fluorescence study disclosed a weakly bound ground state but due to the shape of the potential curves the excited state was found to decay predominantly to the dissociative part of the ground state potential. Hence KMg is shown to be a "Van der Waals" molecule similar to Ca_2 . For laser purposes, however, this species can be considered to be an excimer.

III. RECOMMENDATIONS

The $\text{M}+\text{N}_2\text{O}+\text{CO}$ system shows high promise for development as a chemical laser if the following goals can be accomplished.

1. The $\text{M}+\text{N}_2\text{O}+\text{CO}$ flame as a generator of M^* must be scaled to supply a sufficient M^* flow to power a laser system and the quenching lifetime of the M^* must be long enough to allow efficient M^* generation and subsequent reaction or energy transfer. This may likely occur in supersonic flows. Our studies were subsonic, thus back diffusion mixed reactants and products and lead to rapid M^* quenching by N_2O .
2. A means of extracting the energy in M^* by stimulated emission has to be found. This could occur by supplying a reactant that selectively attacks the $\text{M}(^1\text{S})$ ground state, by energy transfer, or by excimer formation. Our studies have only scratched the surface

here, but already one interesting lead, KMg, has emerged. KMg itself may not be an interesting candidate due to the unfortunate overlapping of the $K_2(B \rightarrow X)$ band system which acts as an absorber. However, this is not an essential feature of this excimer type molecule. Hence, similar bound-free transitions may be found in RbMg or NaMg that do not coincide with absorption by the corresponding alkali dimer. The scaling properties of a chemically pumped excimer laser are very favorable due to the probable short radiative lifetime of the laser transition which imparts a high immunity to quenching once the excited species has been formed. Therefore it should be possible given an adequate supply of Mg^* to pressure scale the chemically pumped excimer laser to a satisfactory gain coefficient.

3. Compatability between the M^* generation scheme and the method of extraction by stimulated emission is required. This may entail some chemical considerations and may favor one means of extraction over another but it is in fact largely an engineering problem in staged injection supersonic reactors.

Therefore, continued experimentation in these areas (M^* scaling, transfer, excimers and $M(^1S)$ scavenging) are highly warranted.

The kind of laser system that might emerge as a result of developments in this field would have the following desirable characteristics.

1. Specific Energy ~400 kJ/lb
2. Wavelength 450-750 nm
3. Plenum Temperature ~1200°C (boiling point of Mg @ 1ATM)

IV. REFERENCES

1. R.W. Field, C.R. Jones, H.P. Broida, J. Chem. Phys. 60 4377 (1974).
2. D.J. Benard, Chem. Phys. Lett. 35, 167 (1975).
3. C.E.H. Bawn, Trans. Far. Soc. 31, 461 (1936).
4. C.P. Fenimore, J.R. Kelso, J. Amer. Chem. Soc. 72, 5045 (1950).
5. A.G. Gaydon, Proc. R. Soc. (London), Ser A 231, 437 (1945).
6. R.W. Pearse and A.G. Gaydon, The Identification of Molecular Spectrum, Wiley and Sons, New York (1963).
7. D.J. Eckstrom, J.R. Baker, J.G. Hawley, Appl. Opt. 16, 2102 (1977).
8. R. Field, private communication.
9. J. Schamps and M. Lefebvre-Brion, J. Chem. Phys. 61, 1652 (1974).
10. M.A. Revelli, B.G. Wicke and D.O. Harris, J. Chem. Phys. 66, 7321 (1977).

APPENDIX

List of Publications (AFOSR 76-2959)

1. D.J. Benard, W.D. Slafer, "Efficient Chemical Production of Metastable Alkaline Earth Atoms," Chem. Phys. Lett. 43, 69-72 (1976).
2. D.J. Benard, W.D. Slafer and J. Hecht, "Chain Reaction Chemiluminescence of Alkaline Earth Catalyzed N_2O -CO Flames," J. Chem. Phys. 66, 1012-1016 (1977).
3. D.J. Benard, W.D. Slafer, "Mechanism of Chemiluminescent Chain Reaction in Mg Catalyzed N_2O -CO Flames," J. Chem. Phys. 66, 1017-1020 (1977).
4. D.J. Benard, W.D. Slafer, J. Hecht and P.H. Lee, "Experimental Investigations and Gain Measurements of Alkaline Earth Catalyzed N_2O +CO Flames," presented at Third Summer Colloquium on Electronic Transition Lasers, Snowmass Village, Colorado (7-10 September 1976). Published in Electronic Transition Lasers II, eds. L.E. Wilson, S.N. Suchard, J.I. Steinfeld, MIT Press, Cambridge, Mass., Chapter II-7 (1977).
5. W.D. Slafer, D.J. Benard and P.H. Lee, "Efficient Chemical Generation of Metastable Alkaline Earth Atoms," presented at Third Summer Colloquium on Electronic Transition Lasers, Snowmass Village, Colorado (7-10 September 1976). Published in Electronic Transition Lasers II, eds. L.E. Wilson, S.N. Suchard, J.I. Steinfeld, MIT Press, Cambridge, Mass., Chapter II-8 (1977).
6. D.J. Benard, P.J. Love and W.D. Slafer, "Energy Transfer and Excimer Formation in a Flowing Afterglow of Mg Metastables," Chem. Phys. Lett., 48, 321-326 (1977). Presented at the IV Conference on Chemical and Molecular Lasers, St. Louis, Missouri, (April 1977).
7. D.J. Benard, W.D. Slafer, P.J. Love and P.H. Lee, "Modulated Transmission Spectroscopy of Gaseous Chemi-Excited Ca and Sr Monoxides," Appl. Optics 16, 2108-2115 (1977).

8. D.J. Benard, W.D. Slafer, "A Mechanistic Study of Potential Chain Reaction Supported Metal Oxide Chemical Lasers, presented at the Annual Electrochemical Society Fall Meeting, Atlanta, Georgia (October 9-14, 1977).
9. W.D. Slafer, D.J. Benard, "Efficient Metal Hydride Arc Source," submitted to Appl. Phys. Lett. (1978).
10. D.J. Benard, W.D. Slafer, "Laser Induced Fluorescence Spectra of KMg," submitted to Chem. Phys. Lett. (1978).